

Physica Status Solidi B: Basic Solid State Physics

A Dyson equation for non-equilibrium Green's functions in the partition-free setting

--Manuscript Draft--

Manuscript Number:	
Full Title:	A Dyson equation for non-equilibrium Green's functions in the partition-free setting
Article Type:	Original Paper
Section/Category:	NonEquilibrium Green's Functions
Keywords:	Non-equilibrium steady-states, Dyson equation
Corresponding Author:	Valeriu Moldoveanu, Ph.D. National Institute of Materials Physics ROMANIA
Additional Information:	
Question	Response
Please submit a plain text version of your cover letter here.	<p>Dear Editors,</p> <p>We would like to submit our paper entitled "A Dyson equation for non-equilibrium Green's functions in the partition-free setting " to be considered for publication as part of the meeting "Progress in Non-Equilibrium Green's function".</p> <p>We consider a small interacting sample coupled to several non-interacting leads. Initially, the system is at thermal equilibrium. At some instant t_0 the system is set into the so called partition-free transport scenario by turning on a bias on the leads. Using the theory of Volterra operators we rigorously formulate a Dyson equation for the retarded Green's function and we establish a closed formula for the associated proper interaction self-energy.</p>
Do you or any of your co-authors have a conflict of interest to declare?	No. The authors declare no conflict of interest.
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	National Institute of Materials Physics
Corresponding Author's Secondary Institution:	
First Author:	Valeriu Moldoveanu, Ph.D.
First Author Secondary Information:	
Order of Authors:	Valeriu Moldoveanu, Ph.D. Claude Alain Pillet, Professor Horia Decebal Cornean
Order of Authors Secondary Information:	
Abstract:	We consider a small interacting sample coupled to several non-interacting leads. Initially, the system is at thermal equilibrium. At some instant t_0 the system is set into the so called partition-free transport scenario by turning on a bias on the leads. Using the theory of Volterra operators we rigorously formulate a Dyson equation for the retarded Green's function and we establish a closed formula for the associated proper interaction self-energy.

A Dyson equation for non-equilibrium Green's functions in the partition-free setting

H. D. Cornean¹, V. Moldoveanu^{*,2}, C.-A. Pillet³

¹ Department of Mathematical Sciences, Aalborg University, Skjernvej 4A, 9220 Aalborg, Denmark

² National Institute of Materials Physics, PO Box MG-7, Bucharest-Magurele, Romania

³ Aix Marseille Univ, Université de Toulon, CNRS, CPT, Marseille, France

Received XXXX, revised XXXX, accepted XXXX

Published online XXXX

Key words: Non-equilibrium Green Functions, Dyson equation, Partition free states

* Corresponding author: e-mail valim@infim.ro

We consider a small interacting sample coupled to several non-interacting leads. Initially, the system is at thermal equilibrium. At some instant t_0 the system is set into the so called partition-free transport scenario by turning on a bias on the leads. Using the theory of Volterra operators we rigorously formulate a Dyson equation for the retarded Green's function and we establish a closed formula for the associated proper interaction self-energy.

Copyright line will be provided by the publisher

1 Introduction The backbone of many-body perturbation theory (MBPT) is the interaction self-energy Σ which appears in the Dyson equation for equilibrium or non-equilibrium Green's function (NEGF). At equilibrium, the structure of Σ is guessed by systematically using Wick's theorem and by analysing the resulting expansion into Feynman diagrams [1]. Approximation schemes (e.g. mean-field approach or RPA) correspond to partial resummation of series of diagrams contributing to Σ .

In the non-equilibrium regime the Gell-Mann and Low Theorem does not hold anymore and writing down statistical averages of time-dependent observables becomes cumbersome. The remedy for these technical difficulties is to combine the chronological T and anti-chronological \bar{T} time-ordering operators into a single operator \mathcal{T}_C which allows an unambiguous book-keeping of time arguments on the two-branch Schwinger-Keldysh contour [2,3]. This construction comes with a price: the non-equilibrium GFs turn to contour-ordered quantities as well and the various identities among them are not easy to recover. At a formal level one assumes the existence of a well-defined self-energy and then the contour-ordered Dyson equation splits via the Langreth rules [4] into the Keldysh equation for the lesser/greater GFs and the Dyson equation for the retarded/advanced GFs (see the textbook [5]).

The existence of a self-energy for the contour-ordered GF is argued by the *formal* analogy between equilibrium and non-equilibrium quantum averages. In fact, a 'complete' self-energy is never written down explicitly. In more recent formulations [6] one starts from the differential equations of motions relating higher n -particle Green-Keldysh functions and then *truncates* the so-called Martin-Schwinger hierarchy [7] to identify various approximate interaction self-energies.

Nowadays, the NEGFs formalism has grown up as a remarkable machinery, being extensively used for modelling quantum transport in mesoscopic systems [8], molecules [9] or even nuclear reactions [10]. Nonetheless, some fundamental theoretical questions were only recently answered by fully exploiting the mathematical structure of the theory and *without* making any approximations. We refer here to: (i) the existence of non-equilibrium steady-state (NESS) in interacting open systems and (ii) the independence of the steady-state quantities from the initial state of the sample [11–15] both in the partitioning [16] and partition free [17,18] settings. We recall here that in the partitioned case the system and the biased leads are initially decoupled.

In our recent work [19] the NEGF formalism for open systems in the partitioning transport setting was rigorously

Copyright line will be provided by the publisher

1 treated in great detail and generality. In particular, we de-
 2 rived the Jauho-Wingreen-Meir formula (JWM) [20] for
 3 the time-dependent current through an interacting sample
 4 by using only real-time quantities.

5 In this short note we are interested in the partition-free
 6 regime which is less studied. We briefly outline a rigorous
 7 formulation of the non-equilibrium Dyson equation for the
 8 retarded Green's function. Mathematical details are kept to
 9 a minimum while focusing on the explicit construction of
 10 a complete interaction self-energy.

11 The content of the paper goes as follows: the model and
 12 the notations are introduced in Section 2, the main result
 13 and its proof are given in Section 3 while Section 4 is left
 14 for conclusions.

15 2 Setting and notation.

16 2.1 Configuration space and Hamiltonians.

17 We assume that a small sample is coupled to M leads. The one-
 18 particle Hilbert space is of tight-binding type and can be
 19 written as $\mathfrak{h} = \mathfrak{h}_S \oplus \mathfrak{h}_R$ where \mathfrak{h}_S is finite dimensional and
 20 $\mathfrak{h}_R = \bigoplus_{\nu=1}^M \mathfrak{h}_\nu$ describes the (finite or not) leads. Particles
 21 can only interact in the sample. One-particle operators are
 22 denoted with lower-case letters and their second quantized
 23 versions will be labeled by capital letters. The one-particle
 24 Hamiltonian of the decoupled system acquires a block-
 25 diagonal structure $h_D = h_S \oplus h_R$ where $h_R = \bigoplus_{\nu=1}^M h_\nu$
 26 is supposed to be bounded. The lead-sample tunnelling
 27 Hamiltonian is defined as:

$$28 \quad h_T = \sum_{\nu=1}^M d_\nu (|f_\nu\rangle\langle g_\nu| + |g_\nu\rangle\langle f_\nu|), \quad (1)$$

29 where ν counts the particle reservoirs, $f_\nu \in \mathfrak{h}_\nu$ and $g_\nu \in$
 30 \mathfrak{h}_S are unit vectors and $d_\nu \in \mathbb{R}$ are coupling constants.
 31 The one-particle Hamiltonian of the fully coupled system
 32 is then $h = h_D + h_T$.

33 We summarize below some useful identities from the
 34 second quantization machinery (see e.g. [21]). The total
 35 Fock space admits a factorization $\mathcal{F} = \mathcal{F}_S \otimes \mathcal{F}_R$. By
 36 $a^\#(f)$ we mean either the creation operator $a^*(f)$ or the
 37 annihilation operator $a(f)$. We have $a^*(\lambda f) = \lambda a^*(f)$
 38 and $a(\lambda f) = \lambda a(f)$. The general form of the canonical
 39 anticommutation relations is:

$$40 \quad \{a(f), a^*(g)\} = \langle f|g\rangle, \quad \{a(f), a(g)\} = 0. \quad (2)$$

41 Here $\langle f|g\rangle$ denotes the scalar product in \mathfrak{h} . Also, $a^\#(f)$ is
 42 bounded on the Fock space and $\|a^\#(f)\| \leq \|f\|$.

43 The interacting, coupled system, and with a potential
 44 bias v_ν on lead ν is described by:

$$45 \quad K_\nu := H + \sum_{\nu=1}^M v_\nu N_\nu + \xi W, \quad (3)$$

46 where N_ν is the particle number operator on lead ν (i.e.,
 47 the second quantization of the orthogonal projection onto

48 \mathfrak{h}_ν), $v := (v_1, \dots, v_M) \in \mathbb{R}^M$ is the bias vector and

$$49 \quad W = \frac{1}{2} \sum_{x,y \in \mathcal{S}} w(x,y) a^*(|x\rangle) a^*(|y\rangle) a(|y\rangle) a(|x\rangle)$$

50 is the second quantization of a two-body potential satisfy-
 51 ing $w(x,y) = w(y,x)$ and $w(x,x) = 0$ for all $x, y \in \mathcal{S}$.
 52 Here $\xi \in \mathbb{R}$ stands for the interaction strength.

53 Assume that the bias is turned on at time $t = 0$. Then
 54 the Heisenberg evolution of an observable A at $t > 0$ is

$$55 \quad \tau_{K_\nu}^t(A) := e^{itK_\nu} A e^{-itK_\nu}, \quad t > 0. \quad (4)$$

56 If h is a single-particle Hamiltonian, the associated
 57 Heisenberg evolution obeys:

$$58 \quad \tau_H^t(a^\#(f)) := e^{itH} a^\#(f) e^{-itH} = a^\#(e^{ith} f), \quad (5)$$

59 and one has

$$60 \quad [H, a^*(f)] = a^*(hf), \quad [H, a(f)] = -a(hf). \quad (6)$$

61 Along the proof of the Dyson equation we shall en-
 62 counter the operators:

$$63 \quad b(f) := i\xi[W, a(f)], \quad b^*(f) := i\xi[W, a^*(f)]. \quad (7)$$

64 These operators vanish if f is supported in the leads.

65 **2.2 The partition-free initial state.** The initial state
 in the partition-free case is a Gibbs state characterized by
 the inverse temperature $\beta > 0$ and the chemical poten-
 tial $\mu \in \mathbb{R}$. It is given by the thermodynamic (i.e., infinite
 leads) limit of the density operator $\rho_{\text{pf}} = Z^{-1} e^{-\beta(K_0 - \mu N)}$
 where $Z = \text{Tr}_{\mathcal{F}} e^{-\beta(K_0 - \mu N)}$. In what follows we briefly
 explain how it is constructed.

The interacting but decoupled and unbiased Hamilto-
 nian is denoted by:

$$66 \quad K_D := H_S + \xi W + H_R = K_0 - H_T.$$

67 The thermodynamic limit of $\rho_D = Z_D^{-1} e^{-\beta(K_D - \mu N)}$
 68 where $Z_D = \text{Tr}_{\mathcal{F}} e^{-\beta(K_D - \mu N)}$ is a tensor product be-
 69 tween a many-body Gibbs state

$$70 \quad \rho_S = \frac{1}{\text{Tr}_{\mathcal{F}_S} e^{-\beta(H_S + \xi W - \mu N_S)}} e^{-\beta(H_S + \xi W - \mu N_S)}$$

71 only acting on the finite dimensional Fock space \mathcal{F}_S , and
 72 M non-interacting (β, μ) Fermi-Dirac quasi-free states
 73 acting on each lead separately, where expectations can
 74 be computed with the usual Wick theorem. This special
 75 factorized initial state is denoted by $\langle \cdot \rangle_{\beta, \mu}$. For exam-
 76 ple, the expectation of a factorized observable of the type
 77 $\mathcal{O} = \mathcal{O}_S \prod_{\nu=1}^M a^*(\tilde{f}_\nu) a(f_\nu)$ where $\tilde{f}_\nu, f_\nu \in \mathfrak{h}_\nu$ is:

$$78 \quad \langle \mathcal{O} \rangle_{\beta, \mu} = \text{Tr}_{\mathcal{F}_S} (\rho_S \mathcal{O}_S) \prod_{\nu=1}^M \langle f_\nu | (\text{Id} + e^{\beta(h_\nu - \mu)})^{-1} \tilde{f}_\nu \rangle.$$

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Its connection with the partition-free state is as follows. Consider the operator $B(\alpha) := e^{-i\alpha K_D} H_T e^{i\alpha K_D}$, $\alpha \in \mathbb{R}$. From (1) and using (5) we see that a generic term entering $B(\alpha)$ is

$$\sum_{\nu} d_{\nu} a^*(e^{-i\alpha h_{\nu}} f_{\nu}) \tau_{H_S + \xi W}^{-\alpha}(a(g_{\nu})).$$

Since h_{ν} is bounded and \mathcal{F}_S is finite dimensional, this expression remains bounded for all complex values of α . Then, the initial value problem

$$\Gamma'(x) = B(ix)\Gamma(x), \quad \Gamma(0) = \text{Id},$$

has a unique solution given by a norm convergent Picard/Dyson/Duhamel iteration, with terms containing products of operators either living in the sample or in the leads. Consequently, writing $e^{-\beta K_0} = e^{-\beta K_D} \Gamma(\beta)$ we obtain an appropriate expression for the thermodynamic limit: \mathcal{O} being an arbitrary bounded physical observable, we have

$$\langle \mathcal{O} \rangle_{\text{pf}} = \frac{\langle \Gamma(\beta) \mathcal{O} \rangle_{\beta, \mu}}{\langle \Gamma(\beta) \rangle_{\beta, \mu}}. \quad (8)$$

2.3 Function spaces and Volterra operators. Let $0 < T < \infty$ be fixed and let $C_0^1([0, T]; \mathfrak{h})$ be the space consisting of time dependent vectors $\phi(t) \in \mathfrak{h}$, $0 \leq t \leq T$, which are continuously differentiable with respect to t , and $\phi(0) = 0$. We also define $C([0, T]; \mathfrak{h})$ to be the space of vectors which are only continuous in t , with no additional condition at $t = 0$. We note that $C([0, T]; \mathfrak{h})$ is a Banach space if we introduce the norm

$$\|\psi\| := \sup_{0 \leq t \leq T} \|\psi(t)\|_{\mathfrak{h}}.$$

We say that an operator A which maps $C([0, T]; \mathfrak{h})$ into itself is a Volterra operator if there exists a constant $C_A < \infty$ such that

$$\|(A\psi)(t)\|_{\mathfrak{h}} \leq C_A \int_0^t \|\psi(t')\|_{\mathfrak{h}} dt', \quad 0 \leq t \leq T.$$

By induction one can prove:

$$\|(A^n \psi)(t)\|_{\mathfrak{h}} \leq C_A \frac{(C_A T)^{n-1}}{(n-1)!} \int_0^t \|\psi(t')\|_{\mathfrak{h}} dt', \quad n \geq 1.$$

In particular, the series $\sum_{n \geq 1} (-1)^n A^n$ converges in operator norm and defines a Volterra operator with a constant less than $C_A e^{T C_A}$. Thus, $(\text{Id} + A)^{-1} = \text{Id} + \sum_{n \geq 1} (-1)^n A^n$ always exists and $A(\text{Id} + A)^{-1}$ is a Volterra operator.

2.4 Retarded NEGF's. Let $\{e_j\}$ be an arbitrary orthonormal basis in \mathfrak{h} . Define the map $G_0 : C([0, T]; \mathfrak{h}) \mapsto C_0^1([0, T]; \mathfrak{h})$ given by:

$$\langle e_j | (G_0 \psi)(t) \rangle := -i \int_0^t \langle e_j | e^{-i(t-t')h_{\nu}} \psi(t') \rangle dt'. \quad (9)$$

One can check that G_0 is invertible and if $\phi \in C_0^1([0, T]; \mathfrak{h})$:

$$(G_0^{-1} \phi)(t) = i \partial_t \phi(t) - h_{\nu} \phi(t) \in C([0, T]; \mathfrak{h}). \quad (10)$$

By definition, the retarded non-equilibrium Green operator in the partition-free setting $G_{\xi} : C([0, T]; \mathfrak{h}) \mapsto C_0^1([0, T]; \mathfrak{h})$ is given by:

$$\langle e_j | (G_{\xi} \psi)(t) \rangle := -i \int_0^t \langle \{ \tau_{K_v}^{t'}(a^*(\psi(t'))), \tau_{K_v}^t(a(e_j)) \} \rangle_{\text{pf}} dt'. \quad (11)$$

Using (5) and (2) we see that G_{ξ} coincides with G_0 when $\xi = 0$. One can show that

$$\|(G_{\xi} \psi)(t)\|_{\mathfrak{h}} \leq 2 \int_0^t \|\psi(t')\|_{\mathfrak{h}} dt', \quad (12)$$

so that G_{ξ} is a Volterra operator. The integral kernel of G_{ξ} is nothing but the more familiar retarded NEGF given by:

$$G_{\xi}^R(e_j, t; e_m, t') := -i \theta(t - t') \langle \{ \tau_{K_v}^{t'}(a^*(e_m)), \tau_{K_v}^t(a(e_j)) \} \rangle_{\text{pf}}, \quad (13)$$

and

$$\langle e_j | (G_{\xi} \psi)(t) \rangle = \sum_m \int_0^t G_{\xi}^R(e_j, t; e_m, t') \langle e_m | \psi(t') \rangle dt'. \quad (14)$$

The advanced NEGF can be defined as:

$$G_{\xi}^A(e_j, t; e_m, t') := -G_{\xi}^R(e_j, t'; e_m, t) = +i \theta(t' - t) \langle \{ \tau_{K_v}^{t'}(a^*(e_m)), \tau_{K_v}^t(a(e_j)) \} \rangle_{\text{pf}}.$$

All properties of the advanced NEGF can be immediately read off from those of the retarded one.

3 Proper self-energy and Dyson equation. Here is the main result of our paper.

Theorem 1. *The bounded linear map $\tilde{\Sigma}_{\xi}$ defined on $C([0, T]; \mathfrak{h})$ by*

$$\langle e_j | (\tilde{\Sigma}_{\xi} \phi)(t) \rangle := -i \int_0^t \langle \{ \tau_{K_v}^{t'}(b^*(\phi(t'))), \tau_{K_v}^t(b(e_j)) \} \rangle_{\text{pf}} dt' + i \langle \tau_{K_v}^t(\{a^*(\phi(t)), b(e_j)\}) \rangle_{\text{pf}}, \quad (15)$$

obeys:

$$G_{\xi} = G_0 + G_0 \tilde{\Sigma}_{\xi} G_0. \quad (16)$$

Moreover, the operator $G_0 \tilde{\Sigma}_{\xi}$ is a Volterra operator, the inverse $(\text{Id} + G_0 \tilde{\Sigma}_{\xi})^{-1}$ exists, and by defining

$$\Sigma_{\xi} := \tilde{\Sigma}_{\xi} (\text{Id} + G_0 \tilde{\Sigma}_{\xi})^{-1} \quad (17)$$

we have:

$$G_\xi = G_0 + G_0 \Sigma_\xi G_\xi. \quad (18)$$

Finally, $G_0 \Sigma_\xi$ is also a Volterra operator and

$$G_\xi = (\text{Id} - G_0 \Sigma_\xi)^{-1} G_0. \quad (19)$$

3.1 Proof: step 1. First we will show that the identity:

$$G_0^{-1} G_\xi = \text{Id} + F_\xi \quad (20)$$

holds on $C([0, T]; \mathfrak{h})$, where the map F_ξ is given by

$$\begin{aligned} \langle e_j | (F_\xi \psi)(t) \rangle := \\ \int_0^t \langle \{ \tau_{K_v}^{t'}(a^*(\psi(t'))), \tau_{K_v}^t(b(e_j)) \} \rangle_{\text{pf}} dt'. \end{aligned} \quad (21)$$

Using (10) and (11) we have:

$$\begin{aligned} \langle e_m | (G_0^{-1} G_\xi \psi)(t) \rangle &= \langle e_m | \psi(t) \rangle \\ &- \sum_j \langle e_m | h_v e_j \rangle \langle e_j | (G_\xi \psi)(t) \rangle \\ &+ \int_0^t \langle \{ \tau_{K_v}^{t'}(a^*(\psi(t'))), \partial_t \tau_{K_v}^t(a(e_m)) \} \rangle_{\text{pf}} dt'. \end{aligned} \quad (22)$$

From the antilinearity of the annihilation operators we get

$$\begin{aligned} \sum_j \langle e_m | h_v e_j \rangle \langle e_j | (G_\xi \psi)(t) \rangle = \\ -i \int_0^t \langle \{ \tau_{K_v}^{t'}(a^*(\psi(t'))), \tau_{K_v}^t(a(h e_m)) \} \rangle_{\text{pf}} dt'. \end{aligned}$$

Also, using (4), (6) and (7) we obtain the identity:

$$\partial_t \tau_{K_v}^t(a(e_m)) = -i \tau_{K_v}^t(h_v a(e_m)) + \tau_{K_v}^t(b(e_m)).$$

After introducing the last two identities into (22) we see that two terms cancel each other and we obtain (21).

3.2 Proof: step 2. The second step consists of showing that F_ξ can be written as $\tilde{\Sigma}_\xi G_0$, with $\tilde{\Sigma}_\xi$ as in (15). In order to identify $\tilde{\Sigma}_\xi$ we compute for every $\phi \in C_0^1([0, T]; \mathfrak{h})$ the quantity (remember that a^* is linear):

$$\begin{aligned} \langle e_j | (F_\xi G_0^{-1} \phi)(t) \rangle \\ = i \int_0^t \langle \{ \tau_{K_v}^{t'}(a^*(\partial_{t'} \phi(t'))), \tau_{K_v}^t(b(e_j)) \} \rangle_{\text{pf}} dt' \\ - \int_0^t \langle \{ \tau_{K_v}^{t'}(a^*(h_v \phi(t'))), \tau_{K_v}^t(b(e_j)) \} \rangle_{\text{pf}} dt'. \end{aligned} \quad (23)$$

Another key identity is:

$$\begin{aligned} \tau_{K_v}^{t'}(a^*(\partial_{t'} \phi(t'))) &= \partial_{t'} \left(\tau_{K_v}^{t'}(a^*(\phi(t'))) \right) \\ &- i \tau_{K_v}^{t'}(a^*(h_v \phi(t'))) - \tau_{K_v}^{t'}(b^*(h_v \phi(t))). \end{aligned}$$

Inserting this identity in (23), integrating by parts with respect to t' and using that $\phi(0) = 0$, we obtain (15).

3.3 Proof: step 3. From the first two steps we derive (16). From (15) and (12) we see that $A = G_0 \tilde{\Sigma}_\xi$ is a Volterra operator for which there exists a T -dependent constant $C < \infty$ such that

$$\|(A\psi)(t)\|_{\mathfrak{h}} \leq C \int_0^t \|\psi(t')\|_{\mathfrak{h}} dt', \quad 0 \leq t \leq T. \quad (24)$$

Then $(\text{Id} + A)^{-1}$ exists and it is given by a norm convergent Neumann series $\sum_{n \geq 0} (-1)^n A^n$, as long as $T < \infty$. We write

$$G_0 = \left(\text{Id} + G_0 \tilde{\Sigma}_\xi \right)^{-1} G_\xi$$

and we can choose Σ_ξ as in (17), which finishes the construction of the proper self-energy.

3.4 Consequences. We list a few remarks concerning our main theorem.

(i) The integral kernel of $\tilde{\Sigma}_\xi$ (see (15)) is given by

$$\begin{aligned} \tilde{\Sigma}_\xi^R(e_j, t; e_m, t') := \\ -i \theta(t - t') \langle \{ \tau_{K_v}^{t'}(b^*(e_m)), \tau_{K_v}^t(b(e_j)) \} \rangle_{\text{pf}} \\ + i \delta(t - t') \langle \tau_{K_v}^t(\{ a^*(e_m), b(e_j) \}) \rangle_{\text{pf}}. \end{aligned} \quad (25)$$

If either e_j or e_m belongs to the leads, then the above matrix element equals zero. The explanation for the first term is that at least one of the two operators $b(e_j)$ and $b^*(e_m)$ defined through (7) would be zero in this case, because the self-interaction W is only supported in the sample, hence it commutes with any observable supported on the leads. For the second term, assume that e_j is from the sample while e_m is from the leads. Then since $b(e_j)$ is a sum of products of three creation/annihilation operators from the sample, it anticommutes with $a^*(e_m)$.

The proper self-energy Σ_ξ has the same support property. One recognizes that $\tilde{\Sigma}_\xi^R(e_j, t; e_m, t')$ is a reducible self-energy. In the diagrammatic language all terms contributing to $\tilde{\Sigma}_\xi^R(e_j, t; e_m, t')$ connect to other diagrams by incoming and outgoing G_0 -lines.

(ii) If both $e_j = x$ and $e_m = y$ are located in the small sample, then from (18) we see that in order to compute $G_\xi^R(x, t; y, t')$ we only need to know the values of G_0 restricted to the small sample (besides Σ_ξ , of course). From (9) we have:

$$G_0^R(x, t; y, t') = -i \theta(t - t') \langle x | e^{-i(t-t')h_v} y \rangle,$$

with $x, y \in \mathcal{S}$. Such matrix elements can be computed from the resolvent $(h_v - z)^{-1}$ restricted to the small sample; we note that via the Feshbach formula, the biased leads appear as a non-local ‘‘dressing’’ potential which perturbs $h_{\mathcal{S}}$, see [14] for details.

At the level of integral kernels, the Dyson equation (18) reads as:

$$G_{\xi}^R(x, t; y, t') = G_0^R(x, t; y, t') + \sum_{u, v \in \mathcal{S}} \int_0^t ds \int_0^s ds' \cdot G_0^R(x, t; u, s) \Sigma_{\xi}^R(u, s; v, s') G_{\xi}^R(v, s'; y, t').$$

(iii) Assume that we can write Σ_{ξ} as $\Sigma_{\text{app}} + \Sigma'$, where Σ_{app} is an approximating Volterra operator. If $G_{\text{app}} = (\text{Id} - G_0 \Sigma_{\text{app}})^{-1} G_0$ is the solution of the approximate Dyson equation $G_{\text{app}} = G_0 + G_0 \Sigma_{\text{app}} G_{\text{app}}$, then we have:

$$G_{\xi} = G_{\text{app}} + G_{\text{app}} \Sigma' G_{\xi}$$

and $G_{\xi} = (\text{Id} - G_{\text{app}} \Sigma')^{-1} G_{\text{app}}$.

(iv) The limit $T \rightarrow \infty$ is a difficult problem. To the best of our knowledge, the only rigorous mathematical results concerning the existence of a steady-state regime in partition free-systems are [14, 15]. Under certain non-resonant conditions and for ξ small enough, one can prove that a quantity like $G_{\xi}^R(e_m, t' + s; e_n, t')$, where $s > 0$ is fixed, will have a limit as $t' \rightarrow \infty$. This is definitely not guaranteed to happen in all cases, not even in non-interacting systems, due to bound states which may produce persistent oscillations.

4 Conclusions We presented a non-perturbative approach to the partition-free transport problem. Starting from the Volterra operator associated to the retarded Green's function we establish its Dyson equation, and we derive closed formulas for the reducible and irreducible self-energies. The proof is rigorous yet elementary in the sense that although the partition-free scenario is a genuine non-equilibrium regime we do not use contour-ordered operators. A Keldysh equation for the lesser Green's function should be established following the same lines of reasoning, with the extra difficulty induced by the fact that in the partition free setting, the small sample is not empty at $t = 0$.

Unravelling the connection between the closed formula (15) and the diagrammatic approach remains an open problem. Although the anti-commutator structure $\langle \tau_{K_v}^t(\{a^*(\phi(t)), b(e_j)\}) \rangle_{\text{pf}}$ in Eq. (15) looks less familiar one can speculate that the systematic application of the Wick theorem should eventually recover various classes of diagrams. A possible approximation in the self-energy would be to replace the interacting propagator $\tau_{K_v}^t(\cdot)$ with the non-interacting one $\tau_{H_v}^t(\cdot)$, where $K_v = H_v + \xi W$. Note however that the application of the Wick theorem is technically challenging due to the extra term $\Gamma(\beta)$ appearing in (8).

Given the fact that the partition-free setting is less studied in the literature, yet more intuitive on physical grounds than the partitioned approach, we hope that our investiga-

tion will trigger more efforts from both the physical and mathematical-physics communities.

Acknowledgements V.M. acknowledges financial support by the CNCS-UEFISCDI Grant PN-III-P4-ID-PCE-2016-0084. H.C. acknowledges financial support by Grant 4181-00042 of the Danish Council for Independent Research | Natural Sciences. C.A.P. acknowledges financial support by the ANR, Grant NON-STOPS (ANR-17-CE40-0006),

References

- [1] A.L. Fetter and J.D. Walecka, *Quantum Theory of Many-Particle Systems*. Dover Publications, New York, 2003.
- [2] J. Schwinger, *J. Math. Phys.* **2**, 407 (1961).
- [3] L.V. Keldysh, *Zh. Eksp. Teor. Fiz.* **47**, 1515 (1964). English translation in *Sov. Phys. JETP* **20**, 1018 (1965).
- [4] D.C. Langreth, in *Linear and Nonlinear Electron Transport in Solids*. J.T. Devreese and V.E. van Doren editors, NATO Advanced Study Institute, Series B: Physics **17**. Plenum Press, New York, 1976.
- [5] H.J.W. Haug and A.-P. Jauho, *Quantum Kinetics in Transport and Optics of Semiconductors*. Springer Series in Solid State Sciences **123**, 2nd Edition. Springer, Berlin, 2007.
- [6] G. Stefanucci and R. van Leeuwen, *Nonequilibrium Many-Body Theory of Quantum System, a Modern Introduction*. Cambridge University Press, Cambridge, 2013.
- [7] M. Scharnke, N. Schlünzen and M. Bonitz, *J. Math. Phys.* **58**, 061903 (2017).
- [8] P. Myöhänen, A. Stan, G. Stefanucci, and R. van Leeuwen, *Phys. Rev. B* **80**, 115107 (2009).
- [9] N.E. Dahlen and R. van Leeuwen, *Phys. Rev. Lett.* **98**, 153004 (2007).
- [10] A. Rios, B. Barker, M. Buchler and P. Danielewicz, *Annals of Physics* **326**, 1274 (2011).
- [11] J. Fröhlich, M. Merkli and D. Ueltschi, *Ann. Henri Poincaré* **4**, 897 (2004).
- [12] W. Aschbacher, V. Jakšić, Y. Pautrat and C.-A. Pillet, in *Open Quantum Systems III. Recent Developments*. S. Attal, A. Joye and C.-A. Pillet editors. Lecture Notes in Mathematics **1882**. Springer, Berlin, 2006.
- [13] V. Jakšić, Y. Ogata and C.-A. Pillet, *Ann. Henri Poincaré* **8**, 1013 (2007).
- [14] H.D. Cornean, V. Moldoveanu and C.-A. Pillet, *Phys. Rev. B* **84**, 075464 (2011).
- [15] H.D. Cornean, V. Moldoveanu and C.-A. Pillet, *Commun. Math. Phys.* **331**, 261 (2014).
- [16] C. Caroli, R. Combescot, P. Nozières and D. Saint-James, *J. Phys. C: Solid State Phys.* **4**, 916 (1971).
- [17] M. Cini, *Phys. Rev. B* **22**, 5887 (1980).
- [18] H.D. Cornean, P. Duclos and R. Purice, *Ann. Henri Poincaré* **13**, 827 (2012).
- [19] H.D. Cornean, V. Moldoveanu and C.-A. Pillet, *Ann. Henri Poincaré* **19**, 411 (2018).
- [20] A.-P. Jauho, N.S. Wingreen and Y. Meir, *Phys. Rev. B* **50**, 5528 (1994).
- [21] Ph.A. Martin and F. Rothen, *Many-Body Problems and Quantum Field Theory – An Introduction*. Springer, Berlin, 2004.